Filed

January 18, 2002

Appl. No. : 09/

09/937,365

AMENDMENTS TO THE CLAIMS

Please amend the Claims as follows. Insertions are shown <u>underlined</u> while deletions are struck through.

- 1-11. (Canceled)
- 12. (Previously presented) The method of claim 24 wherein the administration is selected from the group consisting of orally, intravenously, topically, intramuscularly, intracutaneously, subcutaneously, intraperitoneally, and by aerosolization.
- 13. (Previously presented) The method of claim 12, wherein the administration is orally, admixed with a food product.
- 14. (Previously presented) The method of claim 13, wherein the food product is selected from the group consisting of: juice, soft drinks, teas, powdered soups, jelly, cookies, biscuits, cereal, crackers, candy, breads, noodles, fish paste, chewing gum, ice cream, and chocolate.
- 15. (Previously presented) The method of claim 24 wherein the administration is between one and 4 doses per day.
 - 16-23. (Canceled)
- 24. (Currently amended) A method for treating human pollinosis in a subject, comprising: administering kaempferol-3-glucoside in an <u>effective</u> amount of about 0.48 mg per day per adult to about 3 mg per day per kg of body weight to a subject who suffers from pollinosis.
 - 25-27. (Canceled)
- 28. (New) A method for treating human pollinosis in a subject, comprising: administering kaempferol-3-glucoside in an amount of about 0.48 to about 2 g per day per adult to a subject who suffers from pollinosis.
- 29. (New) A method for treating human pollinosis in a subject, comprising: administering kaempferol-3-glucoside in an amount of about 0.025 to about 3 mg per day per kg of body weight to a subject who suffers from pollinosis.
- 30. (New) The method of Claim 29, wherein said kaempferol-3-glucoside is administered in an amount of about 0.05 to about 1.5 mg per day per kg of body weight to a subject who suffers from pollinosis.